

Annex B. Quick reference to CropWatch indicators, spatial units and methodologies

The following sections give a brief overview of CropWatch indicators and spatial units, along with a description of the CropWatch production estimation methodology. For more information about CropWatch methodologies, visit CropWatch online at www.cropwatch.com.cn.


Agroecological zones for 45 key countries

Overview


228 agroecological zones for the 45 key countries across the globe

Description

45 key agricultural countries are divided into 228 agro-ecological zones based on cropping systems, climatic zones, and topographic conditions. Each country is considered separately. A limited number of regions (e.g., region 001, region 027, and region 127) are not relevant for the crops currently monitored by CropWatch but are included to allow for more complete coverage of the 45 key countries. Some regions are more relevant for rangeland and livestock monitoring, which is also essential for food security.



Asia



- 001. Central region with sparse vegetation
- 002. Dry region
- 003. Mixed dry farming and irrigated cultivation region
- 004. Mixed dry farming and grazing region
- 023. Coastal region
- 024. Ganges plain
- 025. Hills
- 026. Sylhet basin
- 038. Gansu-Xinjiang
- 039. Hainan
- 040. Huang Huihai
- 041. Inner Mongolia
- 042. Loess region
- 043. Lower Yangtze region
- 044. North East China
- 045. Qinghai-Tibet
- 046. Southern China
- 047. South-West China
- 048. China Taiwan
- 090. Java
- 091. Kalimantan and Sulawesi
- 092. Sumatra
- 093. West Papua
- 094. Deccan Plateau
- 095. Eastern coastal region
- 096. Ganges plain
- 097. Assam and north-eastern regions
- 098. Agriculture areas in Rajasthan and Gujarat
- 099. Western coastal region
- 100. North-western dry region
- 101. Western Himalayan region
- 102. Central and Eastern wasteland region
- 103. And Red Sea coastal low hills and plains
- 104. Semi-arid to sub-tropical western and northern hills
- 109. Central non-agriculture region
- 110. South zone
- 111. Eastern plateau and southeastern zone
- 112. Northern zone
- 117. Tonle-sap
- 118. Mekong valley between Tonle-sap and Vietnam border
- 119. Northern plain and northeast
- 120. Southwest Hilly region
- 121. Dry Zone
- 122. Intermediate Zone
- 123. Wet zone
- 132. Central plain
- 133. Delta and southern-coast
- 134. Hills
- 135. Altai
- 136. Gobi Desert
- 137. Hangai Khuvsugul Region
- 138. Selenge-Onon Region
- 139. Central and Eastern Steppe
- 154. Balochistan
- 155. Lower Indus basin in south Punjab and Sind
- 156. Northern highlands
- 157. Northern Punjab
- 158. Forest islands
- 159. Negros and central Visayas islands
- 160. Northern lowlands of Mindanao to western Visayas
- 168. Amur and Primorsky Krai
- 171. Eastern Siberia
- 172. Middle Siberia
- 177. Subarctic region
- 178. Ural and western Volga region
- 179. Western Siberia
- 181. 1st agro-ecological zone
- 182. 1st agro-ecological zone
- 183. Badia
- 184. 2nd agro-ecological zone
- 185. 3rd agro-ecological zone
- 186. 4th agro-ecological zone
- 187. Central double and triple-cropped rice lowlands
- 188. South-eastern horticulture area
- 189. Western and southern hill areas
- 190. Single-cropped rice north-eastern region
- 191. Black Sea region
- 192. Central Anatolia region
- 193. Eastern Anatolia region
- 194. Marmara Aegan Mediterranean lowland region
- 210. Central region with sparse crops
- 211. Eastern hilly cereals zone
- 212. Aral Sea cotton zone
- 213. Central Highlands
- 214. Mekong River Delta
- 215. North Central Coast
- 216. North East
- 217. North West
- 218. Red River Delta
- 219. South Central Coast
- 220. South East

Europe




- 027. Center
- 028. North
- 029. South-west
- 054. East-German lake and Heathland sparse crop area
- 055. Central wheat zone of Saxony and Thuringia
- 056. Wheat zone of Schleswig-Holstein and the Baltic coast
- 057. Mixed wheat and sugarbeets zone of the north-west
- 058. Bavarian Plateau
- 059. Western sparse crop area of the Rhenish massif
- 075. Massid Central dry zone
- 076. Alpes region
- 077. Mediterranean zone
- 078. Northern barley zone
- 079. Maize_barley and livestock zone along the English Channel
- 080. Rapeseed zone of eastern France
- 081. Southwest maize zone
- 082. Mixed maize_barley and rapeseed zone from the Centre to the Atlantic Ocean
- 083. North England, Wales and North Ireland sparse crops area
- 084. Barley area in Scotland
- 085. South English mixed wheat and Barley zone
- 086. Central Hungary
- 087. Puszta
- 088. North Hungary
- 089. Transdanubia
- 105. East coast
- 106. Po Valley
- 107. Islands
- 108. Western Italy
- 161. Central rye and potatoes area
- 162. Northern oats and potatoes areas
- 163. Northern-central wheat and sugarbeet area
- 164. Southern wheat and sugarbeet area
- 165. Central mixed farming and pasture Carpathian hills
- 166. Eastern and southern maize wheat and sugarbeet plains
- 167. Western and central maize wheat and sugarbeet plateau
- 169. Central Russia
- 170. Central black soils area
- 173. Middle Volga
- 174. Northern Caucasus
- 175. Northwest Region including Novgorod
- 176. South Caucasian
- 180. West subarctic region
- 195. Central wheat area
- 196. Eastern Carpathian hills
- 197. Northern wheat area
- 198. Southern wheat and maize area

South America




- 010. Andes
- 011. Chaco
- 012. Mesopotamia
- 013. Humid Pampas
- 014. Pampas hills
- 015. Arid part of Patagonia
- 016. Dry Pampas
- 017. Subtropical highlands
- 030. Amazonas
- 031. Central Savanna
- 032. Coast
- 033. Northeastern mixed forest and farmland
- 034. Mato Grosso
- 035. Nordeste
- 036. Parana basin
- 037. Southern subtropical rangelands

Africa



- 005. Arid Zone
- 006. Central Plateau
- 007. Humid zone
- 008. Semi-Arid Zone
- 009. Sub-humid zone
- 060. Nile Delta and Mediterranean coastal strip
- 061. Nile Valley
- 062. Desert
- 063. Central-northern maize-teff highlands
- 064. Eastern arid area
- 065. Great Rift region
- 066. Northern Arid area
- 067. North-western cereal-root-sesame lowlands
- 068. North-western sesame irrigated lowlands
- 069. North-western semi-arid lowlands
- 070. South-eastern mixed maize zone
- 071. South-eastern Mendebo highlands
- 072. Semi-arid pastoral areas
- 073. South-western coffee-enset highlands
- 074. Western mixed maize zone
- 113. Coast
- 114. Highland agriculture zone
- 115. northern rangelands
- 116. South-west
- 124. Desert
- 125. Sub-humid northern highlands
- 126. Warm semiarid zones
- 127. Warm subhumid zones
- 140. Buzi basin
- 141. Northern high altitude areas
- 142. Low Zambezia River basin
- 143. Northern coast
- 144. Southern region
- 145. Derived Savannah
- 146. Freshwater Swamp Forest
- 147. Guinea Savannah
- 148. Jos Plateau
- 149. Lowland Rainforest
- 150. Mangrove Forest
- 151. Montane Forest
- 152. Sahel Savannah
- 153. Sudan Savannah
- 221. Arid and desert zones
- 222. Humid Cape Fold mountains
- 223. Mediterranean zone
- 224. Dry Highveld and Bushveld maize areas
- 225. Luanguwa Zambezi rift valley
- 226. Northern high rainfall zone
- 227. Central-eastern and southern plateau
- 228. Western semi-arid plain



CropWatch indicators

The CropWatch indicators are designed to assess the condition of crops and the environment in which they grow and develop; the indicators—RAIN (for rainfall), TEMP (temperature), and RADPAR (photosynthetically active radiation, PAR)—are not identical to the weather variables, but instead are value-added indicators computed only over crop growing areas (thus for example excluding deserts and rangelands) and spatially weighted according to the agricultural production potential, with marginal areas receiving less weight than productive ones. The indicators are expressed using the usual physical units (e.g., mm for rainfall) and were thoroughly tested for their coherence over space and time. CWSU are the CropWatch Spatial Units, including MRUs, MPZ, and countries (including first-level administrative districts in select large countries). For all indicators, high values indicate "good" or "positive."

INDICATOR			
BIOMSS			
Biomass accumulation potential			
Crop/ satellite	Grams dry matter/m ² , pixel or CWSU	An estimate of biomass that could potentially be accumulated over the reference period given the prevailing rainfall and temperature conditions.	Biomass is presented as maps by pixels, maps showing average pixels values over CropWatch spatial units (CWSU), or tables giving average values for the CWSU. Values are compared to the average value for the recent fifteen years (2007-2021), with departures expressed in percentage.
CALF			
Cropped arable land and cropped arable land fraction			
Crop/ Satellite	[0,1] number, pixel or CWSU average	The area of cropped arable land as fraction of total (cropped and uncropped) arable land. Whether a pixel is cropped or not is decided based on NDVI twice a month. (For each four-month reporting period, each pixel thus has 8 cropped/uncropped values).	The value shown in tables is the maximum value of the 8 values available for each pixel; maps show an area as cropped if at least one of the 8 observations is categorized as "cropped." Uncropped means that no crops were detected over the whole reporting period. Values are compared to the average value for the last five years (2017-2021), with departures expressed in percentage.
CROPPING INTENSITY			
Cropping intensity Index			
Crop/ Satellite	0, 1, 2, or 3; Number of	Cropping intensity index describes the extent to which arable land is used over	Cropping intensity is presented as maps by pixels or spatial average pixels values for MPZs, 45

INDICATOR			
	crops growing over a year for each pixel	a year. It is the ratio of the total crop area of all planting seasons in a year to the total area of arable land.	countries, and 7 regions for China. Values are compared to the average of the previous five years, with departures expressed in percentage.
NDVI			
Normalized Difference Vegetation Index			
Crop/Satellite	[0.12-0.90] number, pixel or CWSU average	An estimate of the density of living green biomass.	NDVI is shown as average profiles over time at the national level (cropland only) in crop condition development graphs, compared with previous year and recent five-year average (2017-2021), and as spatial patterns compared to the average showing the time profiles, where they occur, and the percentage of pixels concerned by each profile.
RADPAR			
CropWatch indicator for Photosynthetically Active Radiation (PAR), based on pixel based PAR			
Weather /Satellite	W/m ² , CWSU	The spatial average (for a CWSU) of PAR accumulation over agricultural pixels, weighted by the production potential.	RADPAR is shown as the percent departure of the RADPAR value for the reporting period compared to the recent fifteen-year average (2007-2021), per CWSU. For the MPZs, regular PAR is shown as typical time profiles over the spatial unit, with a map showing where the profiles occur and the percentage of pixels concerned by each profile.
RAIN			
CropWatch indicator for rainfall, based on pixel-based rainfall			
Weather / satellite	Liters/m ² , CWSU	The spatial average (for a CWSU) of rainfall accumulation over agricultural pixels, weighted by the production potential.	RAIN is shown as the percent departure of the RAIN value for the reporting period, compared to the recent fifteen-year average (2007-2021), per CWSU. For the MPZs, regular rainfall is shown as typical time profiles over the spatial unit, with a map showing where the profiles occur and the percentage of pixels concerned by each profile.
TEMP			
CropWatch indicator for air temperature, based on pixel-based temperature			
Weather / satellite	°C, CWSU	The spatial average (for a CWSU) of the temperature time average over agricultural pixels, weighted by the production potential.	TEMP is shown as the departure of the average TEMP value (in degrees Centigrade) over the reporting period compared with the average of the recent fifteen years (2007-2021), per CWSU. For the MPZs, regular temperature is illustrated as typical time profiles over the spatial unit, with a map showing where the profiles occur and the percentage of pixels concerned by each profile.
VCIx			
Maximum vegetation condition index			
Crop/Satellite	Number, pixel to CWSU	Vegetation condition of the current season compared with historical data. Values usually are [0, 1], where 0 is "NDVI as bad as the worst recent year" and 1 is "NDVI as good as the best recent year." Values can exceed the range if the current year is the best or the worst.	VCIx is based on NDVI and two VCI values are computed every month. VCIx is the highest VCI value recorded for every pixel over the reporting period. A low value of VCIx means that no VCI value was high over the reporting period. A high value means that at least one VCI value was high. VCI is shown as pixel-based maps and as average value by CWSU.
VHI			
Vegetation health index			

INDICATOR			
Crop/ Satellite	Number, pixel to CWSU	The average of VCI and the temperature condition index (TCI), with TCI defined like VCI but for temperature. VHI is based on the assumption that "high temperature is bad" (due to moisture stress), but ignores the fact that low temperature may be equally "bad" (crops develop and grow slowly, or even suffer from frost).	Low VHI values indicate unusually poor crop condition, but high values, when due to low temperature, may be difficult to interpret. VHI is shown as typical time profiles over Major Production Zones (MPZ), where they occur, and the percentage of pixels concerned by each profile.
VHIn			
Minimum Vegetation health index			
Crop/ Satellite	Number, pixel to CWSU	VHIn is the lowest VHI value for every pixel over the reporting period. Values usually are [0, 100]. Normally, values lower than 35 indicate poor crop condition.	Low VHIn values indicate the occurrence of water stress in the monitoring period, often combined with lower than average rainfall. The spatial/time resolution of CropWatch VHIn is 16km/week for MPZs and 1km/dekad for China.
CPI			
Crop Production Index			
Crop/ Satellite	Number, pixel to CWSU	The average crop production situation for the same period in the past five years was used as a benchmark to make an overall estimate of the current season's agricultural production situation.	Based on the VCIx, CALF, land productivity and area of irrigated and rainfed cropland in the current monitoring period and the same period in the past five years for the spatial unit, a mathematical model proposed by CropWatch is used to calculate the index expressed as a normalized value. A value of 1.0 represents the basic normal crop production situation in the current period for the spatial unit, and the higher the value, the better the crop production situation in the current period. Conversely, the lower the value, the worse the crop production situation for the spatial unit in the current period.

Note: Type is either "Weather" or "Crop"; source specifies if the indicator is obtained from ground data, satellite readings, or a combination; units: in the case of ratios, no unit is used; scale is either pixels or large scale CropWatch spatial units (CWSU). Many indicators are computed for pixels but represented in the CropWatch bulletin at the CWSU scale.

CropWatch spatial units (CWSU)

CropWatch analyses are applied to four kinds of CropWatch spatial units (CWSU): Countries, China, Major Production Zones (MPZ), and global crop Monitoring and Reporting Units (MRU). The tables below summarize the key aspects of each spatial unit and show their relation to each other. For more details about these spatial units and their boundaries, see the CropWatch bulletin online resources.

SPATIAL UNITS	
CHINA	
<i>Overview</i>	<i>Description</i>
Seven monitoring regions	The seven regions in China are agro-economic/agro-ecological regions that together cover the bulk of national maize, rice, wheat, and soybean production. Provinces that are entirely or partially included in one of the monitoring regions are indicated in color on the map below.



Global Monitoring and Reporting Unit (MRU)

Overview	Description
105 agro-ecological/agro-economic units across the world	MRUs are reasonably homogeneous agro-ecological/agro-economic units spanning the globe, selected to capture major variations in worldwide farming and crops patterns while at the same time providing a manageable (limited) number of spatial units to be used as the basis for the analysis of environmental factors affecting crops. Unit numbers and names are shown in the figure below. A limited number of units are not relevant for the crops currently monitored by CropWatch but are included to allow for more complete coverage of global production. Additional information about the MRUs is provided online under www.cropwatch.com.cn .



Production estimation methodology

The main concept of the CropWatch methodology for estimating production is the calculation of current year production based on information about last year's production and the variations in crop yield and cultivated area compared with the previous year. The equation for production estimation is as follows:

$$Production_i = Production_{i-1} * (1 + \Delta Yield_i) * (1 + \Delta Area_i)$$

Where i is the current year, $\Delta Yield_i$ and $\Delta Area_i$ are the variations in crop yield and cultivated area compared with the previous year; the values of $\Delta Yield_i$ and $\Delta Area_i$ can be above or below zero.

For the 45 countries monitored by CropWatch, yield variation for each crop is calibrated against NDVI time series, using the following equation:

$$\Delta Yield_i = f(NDVI_i, NDVI_{i-1})$$

Where $NDVI_i$ and $NDVI_{i-1}$ are taken from the time series of the spatial average of NDVI over the crop specific mask for the current year and the previous year. For NDVI values that correspond to periods after the current monitoring period, average NDVI values of the previous five years are used as an average expectation. $\Delta Yield_i$ is calculated by regression against average or peak NDVI (whichever yields the best regression), considering the crop phenology of each crop for each individual country.

A different method is used for areas. For China, CropWatch combines remote-sensing based estimates of the crop planting proportion (cropped area to arable land) with a crop type proportion (specific type area to total cropped area). The planting proportion is estimated based on an unsupervised classification of high resolution satellite images from HJ-1 CCD and GF-1 images. The crop-type proportion for China is obtained by the GVG instrument from field transects. The area of a specific crop is computed by multiplying farmland area, planting proportion, and crop-type proportion of the crop.

To estimate crop area for wheat, soybean, maize, and rice outside China, CropWatch relies on the regression of crop area against cropped arable land fraction of each individual country (paying due attention to phenology):

$$Area_i = a + b * CALF_i$$

Where, a and b are the coefficients generated by linear regression with area from FAOSTAT or national sources and CALF (Cropped Arable Land Fraction) from CropWatch estimates.